

Openly generated Boolean algebras under FRP

Sakaé Fuchino (湊野 昌)

Kobe University (神戸大学大学院 システム情報学研究科)

`fuchino@diamond.kobe-u.ac.jp`

`http://kurt.scitec.kobe-u.ac.jp/~fuchino/`

(November 9, 2010 (00:53 JST) version)

RIMS Workshop

Interplay between large cardinals and smalls cardinals での講演

於 京都大学数理解析研究所, Kyoto Japan

October 27

This presentation is typeset by p^LA^TE_X with beamer class.

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B (a \leq b \rightarrow \exists c \in f(a) \cap f(b) (a \leq c \leq b))$.

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B (a \leq b \rightarrow \exists c \in f(a) \cap f(b) (a \leq c \leq b))$.

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:

- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B$ ($a \leq b \rightarrow \exists c \in f(a) \cap f(b)$ ($a \leq c \leq b$)).

- ▶ A Boolean algebra B is said to be **openly generated** iff $\{A \in [B]^{\aleph_0} : A \leq_{rc} B\}$ contains a club set ($\subseteq [B]^{\aleph_0}$)
- ▶ $A \leq_{rc} B \Leftrightarrow A$ is a relatively complete subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by a single element (lower projection of b)).

Theorem 1 (S.F., Heindorf, Shapiro, 1994)

For a Boolean algebra B , the following are equivalent:


- (1) B is openly generated;
- (2) $\Vdash_{\mathbb{P}}$ “ B is projective” for any σ -closed \mathbb{P} forcing $|B| = \aleph_1$;
- (3) B has Freese-Nation property. I.e., there is a mapping (Freese-Nation mapping (or FN-mapping)) $f : B \rightarrow [B]^{<\aleph_0}$ s.t. $\forall a, b \in B (a \leq b \rightarrow \exists c \in f(a) \cap f(b) (a \leq c \leq b))$.

A Boolean algebra B is **projective** \Leftrightarrow for any Boolean mapping $\varphi : B \rightarrow A$ and surjective Boolean mapping $\psi : C \rightarrow A$ there is a unique Boolean mapping $\eta : B \rightarrow C$ s.t.

A Boolean algebra B is **projective** \Leftrightarrow for any Boolean mapping $\varphi : B \rightarrow A$ and surjective Boolean mapping $\psi : C \rightarrow A$ there is a unique Boolean mapping $\eta : B \rightarrow C$ s.t.

A Boolean algebra B is **projective** \Leftrightarrow for any Boolean mapping $\varphi : B \rightarrow A$ and surjective Boolean mapping $\psi : C \rightarrow A$ there is a unique Boolean mapping $\eta : B \rightarrow C$ s.t.

A Boolean algebra B is **projective** \Leftrightarrow for any Boolean mapping $\varphi : B \rightarrow A$ and surjective Boolean mapping $\psi : C \rightarrow A$ there is a unique Boolean mapping $\eta : B \rightarrow C$ s.t.



Theorem 2 (S.F. and Q. Feng, 1994)

Assume Axiom R. Then a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective. \square

► B is \aleph_2 -**projective**

$\Leftrightarrow \{C \in [B]^{<\aleph_2} : C \text{ is projective}\}$ contains a club ($\subseteq [B]^{<\aleph_2}$)

Theorem 3 (S.F., 1994)

Assume \square_κ then there is an \aleph_2 -projective but not openly generated Boolean algebra B of cardinality κ^+ . \square

Theorem 2 (S.F. and Q. Feng, 1994)

Assume Axiom R. Then a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective. \square

► B is \aleph_2 -**projective**

$\Leftrightarrow \{C \in [B]^{<\aleph_2} : C \text{ is projective}\}$ contains a club ($\subseteq [B]^{<\aleph_2}$)

Theorem 3 (S.F., 1994)

Assume \square_κ then there is an \aleph_2 -projective but not openly generated Boolean algebra B of cardinality κ^+ . \square

Theorem 2 (S.F. and Q. Feng, 1994)

Assume Axiom R. Then a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective. \square

► B is \aleph_2 -projective

$\Leftrightarrow \{C \in [B]^{<\aleph_2} : C \text{ is projective}\}$ contains a club ($\subseteq [B]^{<\aleph_2}$)

Theorem 3 (S.F., 1994)

Assume \square_κ then there is an \aleph_2 -projective but not openly generated Boolean algebra B of cardinality κ^+ . \square

Theorem 2 (S.F. and Q. Feng, 1994)

Assume Axiom R. Then a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective. \square

► B is \aleph_2 -**projective**

$\Leftrightarrow \{C \in [B]^{<\aleph_2} : C \text{ is projective}\}$ contains a club ($\subseteq [B]^{<\aleph_2}$)

Theorem 3 (S.F., 1994)

Assume \square_κ then there is an \aleph_2 -projective but not openly generated Boolean algebra B of cardinality κ^+ . \square

Theorem 2 (S.F. and Q. Feng, 1994)

Assume Axiom R. Then a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective. \square

► B is \aleph_2 -**projective**

$\Leftrightarrow \{C \in [B]^{<\aleph_2} : C \text{ is projective}\}$ contains a club ($\subseteq [B]^{<\aleph_2}$)

Theorem 3 (S.F., 1994)

Assume \square_κ then there is an \aleph_2 -projective but not openly generated Boolean algebra B of cardinality κ^+ . \square

Main Theorem 4 (S.F. and A. Rinot, to appear)

The assertion of Theorem 2:

a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective

is equivalent to FRP over ZFC.

► **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in S \cap I$;
- for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \upharpoonright \{\xi^*\}$ is stationary in $\text{sup}(I)$.

Main Theorem 4 (S.F. and A. Rinot, to appear)

The assertion of Theorem 2:

a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective

is equivalent to FRP over ZFC.

► **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in S \cap I$;
- for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \parallel \{\xi^*\}$ is stationary in $\text{sup}(I)$.

Main Theorem 4 (S.F. and A. Rinot, to appear)

The assertion of Theorem 2:

a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective

is equivalent to FRP over ZFC.

► **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in S \cap I$;
- for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \parallel \{\xi^*\}$ is stationary in $\text{sup}(I)$.

Main Theorem 4 (S.F. and A. Rinot, to appear)

The assertion of Theorem 2:

a Boolean algebra B is openly generated if and only if B is \aleph_2 -projective

is equivalent to FRP over ZFC.

► **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in S \cap I$;
- for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \parallel \{\xi^*\}$ is stationary in $\text{sup}(I)$.

- ▶ **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- ▶ $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- ▶ for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \parallel \{\xi^*\}$ is stationary in $\text{sup}(I)$.

- ▶ FRP is introduced and studied in [S.F., Juhász, Soukup, Szentmiklóssy and Usuba 2010].

- ▶ $\text{MM} \Rightarrow \text{MA}^+(\sigma\text{-closed}) \Rightarrow \text{Axiom R} \Rightarrow \text{RP} \not\stackrel{\text{L}}{\Rightarrow} \text{FRP} \not\stackrel{\text{L}}{\Rightarrow} \text{ORP}$

([S.F. et al. 2010] and [S.F., Sakai, Soukup and Usuba 201?])

- ▶ **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- ▶ $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- ▶ for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \parallel \{\xi^*\}$ is stationary in $\text{sup}(I)$.

- ▶ FRP is introduced and studied in [S.F., Juhász, Soukup, Szentmiklóssy and Usuba 2010].

- ▶ $\text{MM} \Rightarrow \text{MA}^+(\sigma\text{-closed}) \Rightarrow \text{Axiom R} \Rightarrow \text{RP} \not\Leftarrow \text{FRP} \not\Leftarrow \text{ORP}$

([S.F. et al. 2010] and [S.F., Sakai, Soukup and Usuba 201?])

- ▶ **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- ▶ $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- ▶ for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \parallel \{\xi^*\}$ is stationary in $\text{sup}(I)$.

- ▶ FRP is introduced and studied in [S.F., Juhász, Soukup, Szentmiklóssy and Usuba 2010].

- ▶ $\text{MM} \Rightarrow \text{MA}^+(\sigma\text{-closed}) \Rightarrow \text{Axiom R} \Rightarrow \text{RP} \not\stackrel{\Leftarrow}{\Rightarrow} \text{FRP} \not\stackrel{\Leftarrow}{\Rightarrow} \text{ORP}$

([S.F. et al. 2010] and [S.F., Sakai, Soukup and Usuba 201?])

- ▶ **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- ▶ $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- ▶ for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \upharpoonright \{\xi^*\}$ is stationary in $\text{sup}(I)$.

- ▶ FRP is introduced and studied in [S.F., Juhász, Soukup, Szentmiklóssy and Usuba 2010].

- ▶ $\text{MM} \Rightarrow \text{MA}^+(\sigma\text{-closed}) \Rightarrow \text{Axiom R} \Rightarrow \text{RP} \not\stackrel{\Leftarrow}{\Rightarrow} \text{FRP} \not\stackrel{\Leftarrow}{\Rightarrow} \text{ORP}$

([S.F. et al. 2010] and [S.F., Sakai, Soukup and Usuba 201?])

► **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \upharpoonright \{\xi^*\}$ is stationary in $\text{sup}(I)$.

Lemma 5

FRP is equivalent to its following variant:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there are stationarily many $I \in [\kappa]^{\aleph_1}$ such that

- $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- $\{x \in [I]^{\aleph_0} : g(\text{sup}(x)) \cap \text{sup}(x) \subseteq x\}$ is stationary.

- ▶ **FRP** (Fodor-type Reflection Principle) is the following assertion:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there is $I \in [\kappa]^{\aleph_1}$ such that

- ▶ $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- ▶ for any $f : S \cap I \rightarrow \kappa$ s.t. $f(\alpha) \in g(\alpha) \cap \alpha$ for all $\alpha \in S \cap I$, there is $\xi^* < \kappa$ s.t. $f^{-1} \upharpoonright \{\xi^*\}$ is stationary in $\text{sup}(I)$.

Lemma 5

FRP is equivalent to its following variant:

For any regular cardinal $\kappa > \aleph_1$, any stationary $S \subseteq E_\omega^\kappa$ and $g : S \rightarrow [\kappa]^{\aleph_0}$ there are stationarily many $I \in [\kappa]^{\aleph_1}$ such that

- ▶ $\text{cf}(I) = \omega_1$; $g(\alpha) \subseteq I$ for all $\alpha \in I \cap S$;
- ▶ $\{x \in [I]^{\aleph_0} : g(\text{sup}(x)) \cap \text{sup}(x) \subseteq x\}$ is stationary.

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \iff A$ is a sigma subalgebra of B

$\iff A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \iff A$ is a sigma subalgebra of B

$\iff A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \iff A$ is a sigma subalgebra of B

$\iff A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \Leftrightarrow A$ is a sigma subalgebra of B

$\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \iff A$ is a sigma subalgebra of B
 $\iff A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \iff A$ is a sigma subalgebra of B

$\iff A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

► SSH (Shelah's Strong Hypothesis) is equivalent to the assertion that $\text{cf}([\lambda]^{\aleph_0}, \subseteq) = \lambda^+$ for all regular cardinal with $\text{cf}(\lambda) = \omega$.

Theorem 7 (S.F.)

Suppose that SSH holds. Then every \aleph_2 -projective Boolean algebras B have a filtration $\langle B_\alpha : \alpha < \kappa \rangle$ for $\kappa = \text{cf}(|B|)$ s.t. $B_{\alpha+1}$ is \aleph_2 -projective and $B_{\alpha+1} \leq_\sigma B$ for all $\alpha < \kappa$. In particular, we also have $B_\alpha \leq_\sigma B$ for all limit $\alpha < \kappa$ of countable cofinality.

► $A \leq_\sigma B \Leftrightarrow A$ is a sigma subalgebra of B
 $\Leftrightarrow A$ is a subalgebra of B and $\forall b \in B$ (the ideal $A \upharpoonright b$ is generated by countable subset of $A \upharpoonright b$).

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all B of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Sketch of the proof of Main Theorem (1/4)

Openly generated Bas (9/14)

▶ “ B is openly generated $\Rightarrow B$ is \aleph_2 -projective” is easy and provable in ZFC.

▶ $\text{FRP} \Rightarrow (B \text{ is } \aleph_2\text{-projective} \Rightarrow B \text{ is openly generated})$ (*)

By induction on $|B|$.

▷ If $|B| \leq \aleph_1$ this is trivial.

▷ Suppose (*) is true for all Ba of cardinality $< |B| = \lambda$.

Case I: λ is singular: a proof similar to that of Shelah’s Singular Compactness Theorem will do.

Case II: λ is regular:

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated!). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\neg\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated !). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\neg\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated !). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Case II: λ is regular:

Suppose that B is an \aleph_2 -projective non openly generated Boolean algebra with $|B| = \lambda$.

By Theorem 7, there is a filtration $\langle B_\alpha : \alpha < \lambda \rangle$ of B s.t. $B_\alpha \leq_\sigma B$ for all $\alpha \in \lambda \setminus E_{>\omega}^k$. The filtration can be chosen so that each B_α is \aleph_2 -projective. By the induction hypothesis, all B_α , $\alpha < \lambda$ are openly generated.

$E = \{\alpha \in E_\omega^\lambda : B_\alpha \leq_{\text{rc}} B\}$ is stationary (otherwise we could conclude that B is openly generated !). Without loss of generality $B_\alpha = \alpha$ for all $\alpha \in E$.

For $\alpha \in E$, there is $\eta_\alpha \in \lambda \setminus \alpha$ s.t. $B_\alpha \upharpoonright \eta_\alpha$ is countably generated by some $c_\alpha \in [B_\alpha \upharpoonright \eta_\alpha]^{\aleph_0}$ but not by any single element. Let $g : E \rightarrow [\lambda]^{\aleph_0}$ be defined by $g(\alpha) = c_\alpha \cup \{\eta_\alpha\}$ for $\alpha \in E$.

Sketch of the proof of Main Theorem (3/4)

Openly generated Bas (11/14)

By the characterization (Lemma 5) of FRP, there is $I \in [\lambda]^{\aleph_1}$ s.t. I is prjective as a subalgebra of B , $\text{cf}(I) = \omega_1$, I is closed w.r.t. g and $\{x \in [I]^{\aleph_0} : g(\text{sup}(x)) \cap \text{sup}(x) \subseteq x\}$ is stationary.

From the last condition it follows that I is not projective which is a contradiction.

Sketch of the proof of Main Theorem (3/4)

Openly generated Bas (11/14)

By the characterization (Lemma 5) of FRP, there is $I \in [\lambda]^{\aleph_1}$ s.t. I is projective as a subalgebra of B , $\text{cf}(I) = \omega_1$, I is closed w.r.t. g and $\{x \in [I]^{\aleph_0} : g(\text{sup}(x)) \cap \text{sup}(x) \subseteq x\}$ is stationary.

From the last condition it follows that I is not projective which is a contradiction.

Sketch of the proof of Main Theorem (3/4)

Openly generated Bas (11/14)

By the characterization (Lemma 5) of FRP, there is $I \in [\lambda]^{\aleph_1}$ s.t. I is projective as a subalgebra of B , $\text{cf}(I) = \omega_1$, I is closed w.r.t. g and $\{x \in [I]^{\aleph_0} : g(\text{sup}(x)) \cap \text{sup}(x) \subseteq x\}$ is stationary.

From the last condition it follows that I is not projective which is a contradiction.

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

► (B is \aleph_2 -projective $\Leftrightarrow B$ is openly generated) \Rightarrow FRP:

Assume \neg FRP. Then there is a regular $\kappa > \aleph_1$, stationary $S \subseteq E_\omega^\kappa$ and a ladder system $g : S \rightarrow [\kappa]^{\aleph_0}$ s.t., for any $\alpha < \kappa$, there is a regressive $f : S \cap \alpha \rightarrow \alpha$ s.t. $\{g(\beta) \setminus f(\beta) : \beta \in S \cap \alpha\}$ is pairwise disjoint ([S.F., Sakai, Soukup and Usuba 201?]).

Let $D = \kappa \setminus \text{Lim}$. Without loss of generality, $g(\alpha) \subseteq D$ for all $\alpha \in S$. Let $X = \{c_\alpha : \alpha \in S \cup D\}$.

Let $<_B$ be a binary relation on X defined by

$$c_\alpha <_B c_\beta \Leftrightarrow \alpha \in D \wedge \beta \in S \wedge \alpha \in g(\beta).$$

Let B be the Ba generated over X freely except $<_B$.

► Then B is \aleph_2 -projective but not openly generated.

□ (Main Theorem)

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

Theorem 6 (A. Rinot, independently: T. Usuba)

FRP *implies* SSH.

(Very Rough) Sketch of the Proof:

Suppose that SSH does not hold. Then there is a better scale $\langle \langle \lambda_i : i < \omega \rangle, \langle f_\alpha : \alpha < \lambda^+ \rangle \rangle$ for a cardinal λ with $\text{cf}(\lambda) = \omega$ ([Shelah: Card. Arith.]).

Let $\varphi : {}^\omega \lambda \rightarrow \lambda$ be a 1-1 mapping, $E = E_\omega^{\lambda^+} \setminus \lambda$ and let $g : E \rightarrow [\lambda^+]^{\aleph_0}$ be s.t. $g(\alpha) = \{\varphi(f_\alpha \upharpoonright n) : n \in \omega\}$.

Then g together with E is a counterexample to $\text{FRP}(\lambda^+)$. □

ご静聴ありがとうございました!

Openly generated Bas (14/14)

Thank you for your attention!



M.C. Escher, "Rippled Surface" (1950)

These slides and their printer friendly version are linked to:
<http://kurt.scitec.kobe-u.ac.jp/~fuchino/>

ご静聴ありがとうございました!

Openly generated Bas (14/14)

Thank you for your attention!



M.C. Escher, "Rippled Surface" (1950)

These slides and their printer friendly version are linked to:
<http://kurt.scitec.kobe-u.ac.jp/~fuchino/>