

Rado's conjecture and reflection principles compatible with MM

Sakaé Fuchino (渕野 昌)

Graduate School of System Informatics
Kobe University

(神戸大学大学院 システム情報学研究科)

<http://kurt.scitec.kobe-u.ac.jp/~fuchino/>

日本数学会 2014 年年会

(21. März 2014 (16:14 JST) version)

March 16, 2014, 於 学習院大学

This presentation is typeset by p^AT_EX with beamer class.

- ▶ A tree T is **special** if there are T_i , $i \in \omega$ s.t. all of T_i 's are pairwise incomparable (antichains) and $T = \bigcup_{i \in \omega} T_i$.

- ▶ Rado's Conjecture (RC):

RC: Any tree T is special if and only if all subtrees of T of cardinality \aleph_1 are special.

- ▷ (S. Todorčević) Rado's Conjecture implies the non-existence of Kurepa trees. In particular RC does not hold under $V = L$.
- ▷ (S. Todorčević) If κ is strongly compact and $\mathbb{P} = \text{Col}(\omega_1, <\kappa)$, then we have $\Vdash_{\mathbb{P}}$ "Rado's Conjecture".
- ▶ Reflection Principle (RP):

RP: For any regular cardinal $\kappa > \aleph_1$ and stationary $\mathcal{S} \subseteq [\kappa]^{\aleph_0}$, there is $I \in [\kappa]^{\aleph_1}$ s.t. (1) $\omega_1 \subseteq I$, (2) $\text{cf}(I) = \text{cf} \sup(I) = \omega_1$ and (3) $\mathcal{S} \cap [I]^{\aleph_0}$ is stationary in $[I]^{\aleph_0}$.

Rado's Conjecture and the Reflection Principle RP (2/2)_{RC-RP} (3/15)

RC: Any tree T is special if and only if all subtrees of T of cardinality \aleph_1 are special.

RP: For any cardinal regular cardinal $\kappa > \aleph_1$ and stationary $\mathcal{S} \subseteq [\kappa]^{\aleph_0}$, there is $I \in [\kappa]^{\aleph_1}$ s.t. (1) $\omega_1 \subseteq I$, (2) $\text{cf}(I) = \text{cf} \sup(I) = \omega_1$ and (3) $\mathcal{S} \cap [I]^{\aleph_0}$ is stationary in $[I]^{\aleph_0}$.

▷ (S. Todorčević, ??) If $V = L$ we have $\neg\text{RC}$ and $\neg\text{RP}$

▷ (S. Todorčević, Foreman-Magidor-Shelah (?))

If κ is supercompact and $\mathbb{P} = \text{Col}(\omega_1, < \kappa)$, then we have

$\Vdash_{\mathbb{P}} \text{“RC} \wedge \text{RP”}$.

- ▶ RC and RP have many common consequences:
 - ▷ $2^{\aleph_0} \leq \aleph_2$;
 - ▷ $\neg \square_\kappa$ for all κ ;
 - ▷ Singular Cardinal Hypothesis;
 - ▷ Chang's Conjecture (and hence the non existence of Kurepa trees);
 - ▷ Ordinal Stationarity Reflection: For any regular $\kappa > \omega_1$ and stationary $S \subseteq S_\kappa^\omega = \{\alpha \in \kappa : \text{cf}(\alpha) = \omega\}$ there is an $\xi \in S_\kappa^{\omega_1}$ s.t. $S \cap \xi$ is stationary in ξ ;
 - ▷ ... (will be discussed later)

Are RC and RP perhaps the same principle? Or at least isn't it so that one of them can be derived from the other?

Neither!!

- ▶ (S. Todorčević) RC implies the negation of Martin's Axiom for \aleph_1 dense sets: RP follows from Martin's Maximum.
- ▷ Hence (under a supercompact cardinal) $\neg\text{RC} + \text{RP}$ is consistent. (MM implies the combination!)
- ▶ (酒井拓史 (H. Sakai)) (Under a supercompact cardinal) it is consistent that there is a strongly compact cardinal κ s.t., for $\mathbb{P} = \text{Col}(\omega_1, <\kappa)$, we have $\Vdash_{\mathbb{P}} \neg\text{RP}$.
- ▷ Hence (under a supercompact cardinal) $\text{RC} + \neg\text{RP}$ is consistent.

Theorem 1. (S. Todorćević) MA_{\aleph_1} implies the negation of RC.

Proof. Let $T = \{t : t \text{ is an increasing sequence in } \mathbb{R} (\ell(t) < \omega_1)\}$ with the ordering $t <_T t' \Leftrightarrow t'$ is an endextension of t .

Then

- ▷ T is a non-special tree.
- ▷ Under MA_{\aleph_1} all subtrees of cardinality \aleph_1 are special by a theorem of Baumgartner-Malitz-Reinhard:
 - (MA_{\aleph_1}) all trees of cardinality \aleph_1 without uncountable chain are special. □

Theorem 2. (1) (P. Doebler, 2013) RC implies the Semi-Stationary Reflection Principle (SSR).

(2) (S.F., H. Sakai, V. Torres and T. Usuba, ∞) RC implies the Fodor-type Reflection Principle (FRP).

Since it is already known that RP implies both SSR and FRP, we obtain the diagram:



SSR implies:

FRP implies:


- ▷ $2^{\aleph_0} \leq \aleph_2$
- ▷ $\neg \square_\kappa$ for all κ
- ▷ Singular Cardinal Hypothesis
- ▷ Chang's Conjecture (and hence the non existence of Kurepa trees)
- ▷ Ordinal Stationarity Reflection: For any regular $\kappa > \omega_1$ and stationary $S \subseteq S_\kappa^\omega = \{\alpha \in \kappa : \text{cf}(\alpha) = \omega\}$ there is an $\xi \in S_\kappa^{\omega_1}$ s.t. $S \cap \xi$ is stationary in ξ
- ▷ ... (will be discussed later)

- ▶ FRP is known to be equivalent (over ZFC) to many “mathematical” reflection theorems such as:
 - ▷ Any locally countably compact space X is metrizable if all subspaces of X of cardinality $\leq \aleph_1$ are metrizable.
 - ▷ Any T_1 -space with point countable base is left separated if every subspaces of X of cardinality $\leq \aleph_1$ are left separated.
 - ▷ Any graph G is of countable coloring number if all subgraphs Y of X of cardinality $\leq \aleph_1$ are of countable coloring number.
 - ▷ ...

- For a regular cardinal $\kappa > \aleph_1$

$CC^\downarrow(\kappa)$: For any sufficiently large regular θ and an well-ordering \sqsubset on $\mathcal{H}(\theta)$, if $\kappa \in M \prec \langle \mathcal{H}(\theta), \in, \sqsubset \rangle$ is countable then, for any $\alpha < \kappa$, there is a countable $M \prec M^* \prec \langle \mathcal{H}(\theta), \in, \sqsubset \rangle$ s.t., letting $\alpha^* = \inf((\kappa \cap M^* \setminus \sup(\kappa \cap M)))$, we have $\alpha^* > \alpha$ and $\text{cf}(\alpha^*) = \omega_1$.

$CC^*(\kappa)$: (P. Doebler) For any sufficiently large regular θ and an well-ordering \sqsubset on $\mathcal{H}(\theta)$, if $\kappa \in M \prec \langle \mathcal{H}(\theta), \in, \sqsubset \rangle$ is countable then, for any $a \in [\kappa]^{\aleph_1}$, there is a countable $M \prec M^* \prec \langle \mathcal{H}(\theta), \in, \sqsubset \rangle$ s.t., $M \cap \omega_1 = M^* \cap \omega_1$ and there is a $b \in [\kappa]^{\aleph_1} \cap M^*$ s.t. $a \subseteq b$.



- ▶ $CC^\downarrow(\kappa)$ and $CC^*(\kappa)$ suggest the following natural generalization of the both of the principles:

$CC^{\downarrow\downarrow}(\kappa)$: For any sufficiently large regular θ and a well-ordering \sqsubset on $\mathcal{H}(\theta)$, if $M \prec \langle \mathcal{H}(\theta), \in, \sqsubset \rangle$ is countable with $\kappa \in M$ then, for any $a \in [\kappa]^{\aleph_1}$, there are $b \in [\kappa]^{\aleph_1}$ and countable $M \prec M^* \prec \mathcal{M}$ s.t. $a \subseteq b$, $b \in M^*$ and $b \cap M^* = b \cap M$.

- ▶ The principle $CC^{\downarrow\downarrow}(\kappa)$ for all $\kappa > \aleph_1$ clearly implies both FRP and SSR. Unfortunately, this principle is a strengthening of RP and hence RC does not imply this principle:


Theorem 3. (S.F., 薄葉季路 (T. Usuba)) For all κ T.f.a.e:

(a) $CC^{\downarrow\downarrow}(\kappa)$ for all $\kappa > \aleph_1$.


(b) For all κ and any stationary $S \subseteq [\kappa]^{\aleph_0}$, for all sufficiently large regular θ and well ordering \sqsubset on $\mathcal{H}(\theta)$, there is an $M \prec \langle \mathcal{H}(\theta), \in, \sqsubset \rangle$ of cardinality \aleph_1 , s.t. $S \cap M$ is stationary in $[\kappa \cap M]^{\aleph_0}$.

Possible Principle(s) unifying FRP and SSR(2/2)


RC-RP (12/15)



- ▶ The model obtained by Levy collapsing (by $\mathbb{P} = \text{Col}(\omega_1, <\kappa)$) a supercompact cardinal can be seen as quite “canonical”.
- ▶ Game Reflection Principle GRP^+ of B. Koenig captures many features of this model. In particular GRP^+ implies RC, RP as well as CH.



- ▶ Under $\neg\text{CH}$, “the canonical model $\models \text{MM}$ ” does not satisfy RC.
- ▶ Mitchel’s model constructed starting from a supercompact cardinal satisfies both RC and RP under $2^{\aleph_0} = \aleph_2$. This model is however “less canonical”.
- ▷ Is there an axiom which captures a good deal of the characteristics of the Mitchel model?





終